



# Order information for Aermony RespiClick™ (wholesale)

## About Aermony RespiClick™







232 mcg

 $\rightarrow$  Therapeutic class

Corticosteroid for oral inhalation

 $\rightarrow \ \, \text{Active ingredient}$ 

Fluticasone propionate inhalation powder, Mfr. Std.

ightarrow Clinically relevant nonmedicinal ingredients

Lactose monohydrate (contains milk protein)

→ Format

Dry powder for inhalation

Product name	Item code	UPC	DIN/NPN	McKesson	K&F	Familiprix	McMahon	PJC	NuQuest	Imperial BC	Imperial AB
Aermony RespiClick™ 55 mcg 60 dose trade	83450-3035	068510178893	02467895	154825	163715	143259	10124260	751175	AERM055TEV	998893	876383
Aermony RespiClick™ 113 mcg 60 dose trade	83451-3035	068510178909	02467909	154821	163716	1142951	10124271	751178	AERM113TEV	998909	876391
Aermony RespiClick™ 232 mcg 60 dose trade	83454-3035	068510178930	02467917	154828	1163717	142957	10124272	751205	AERM232TEV	998930	876441

## **Storage instructions**

Store Aermony RespiClick<sup>™</sup> at room temperature (between 15 °C and 25 °C) in a dry place; excursions permitted from 15 °C to 30 °C. Avoid exposure to extreme heat, cold, or humidity. Aermony RespiClick<sup>™</sup> should be stored inside the unopened moisture-protective foil pouch and only removed from the pouch immediately before initial use. Discard Aermony RespiClick<sup>™</sup> 30 days after opening the foil pouch or when the counter reads "0," whichever comes first. The inhaler is not reusable. Do not attempt to take the inhaler apart. Keep out of reach of children.

Aermony RespiClick™ is indicated for the maintenance treatment of steroid-responsive bronchial asthma as prophylactic therapy in patients 12 years of age and older.

Consult the Product Monograph at https://health-products.canada.ca/dpd-bdpp/index-eng.jsp for contraindications, warnings, precautions, adverse reactions, interactions, dosing, and conditions of clinical use. The Product Monograph is also available by calling us at 1-855-514-8382.

**Reference:** Aermony RespiClick™ Product Monograph. Teva Canada. December 14, 2018.







fluticasone propionate inhalation powder

# Overview of how to use Aermony RespiClick®

#### INSTRUCTIONS FOR PATIENTS

For more information and an instructional video, visit

## www.respiclick.ca

Scan this QR code or type the URL into your browser.



Log in with the DIN on the outside of your Aermony RespiClick® package or with this number: 02467895.

# To use: Open. Inhale. Close.



a "click"—once the cap clicks, one dose is ready to be inhaled.

Make sure you are holding the inhaler upright. Before going to step 2, hold the inhaler away from your mouth and breathe out through your mouth as much air as you can and as is comfortable.

- Never breathe out into the inhaler mouthpiece.
- · Do not open the green cap unless you are taking a dose.



# Inhale.

INHALE quickly and deeply through the mouth, until lungs feel completely full of air.

Your lips should form a good seal around the mouthpiece, but the vent above the mouthpiece should not be blocked by the lips or fingers. Remove the inhaler from your mouth and hold your breath for about 10 seconds or for as long as you comfortably can.

 Do not take an extra dose from the inhaler even if you do not taste or feel the medicine.



be ready for the next dose.

**Rinse your mouth with water after taking your dose.** Spit out the water. Do not swallow it.

Throw away your inhaler 30 days after you have opened the foil pouch or when the counter reads "0," whichever comes first.

Please read the medication leaflet that came with your inhaler for more information about how to take Aermony RespiClick®.







Aermony RespiClick® is a registered trademark of Ivax International B.V., a wholly-owned subsidiary of Teva Pharmaceutical Industries Ltd., and is used under licence.

TEVA and the design version thereof are registered trademarks

of Teva Pharmaceutical Industries Ltd. and are used under licence.

© 2021 Teva Canada Innovation G.P. – S.E.N.C., Montreal, Quebec H2Z 1S8